

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

Agronomy & Horticulture -- Faculty Publications

Agronomy and Horticulture Department

7-2001

Symposium 15. Photosynthesis and respiration Paper S15-022: Lack of Genetic Adaptation and Its Effect on Dark Respiration

F. Zavala-Garcia

Facultad de Agronomia. UANL. Monterrey, N.L. Mexico. Fax: (52) 8248-0101

Jerry Eastin

University of Nebraska-Lincoln, jeastin1@unl.edu

H. Gamez-Gonzalez

Facultad de Ciencias Biologicas. UANL. Monterrey, N.L. Mexico

Follow this and additional works at: <https://digitalcommons.unl.edu/agronomyfacpub>

 Part of the [Plant Sciences Commons](#)

Zavala-Garcia, F.; Eastin, Jerry; and Gamez-Gonzalez, H., "Symposium 15. Photosynthesis and respiration Paper S15-022: Lack of Genetic Adaptation and Its Effect on Dark Respiration" (2001). *Agronomy & Horticulture -- Faculty Publications*. 314.

<https://digitalcommons.unl.edu/agronomyfacpub/314>

This Article is brought to you for free and open access by the Agronomy and Horticulture Department at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Agronomy & Horticulture -- Faculty Publications by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

Symposium 15. Photosynthesis and respiration

Paper S15-022:

Lack of genetic adaptation and its effect on dark respiration

F. Zavala-Garcia,¹ J. D. Eastin,² and H. Gamez-Gonzalez³

1 Facultad de Agronomia. UANL. Monterrey, N.L. Mexico. Fax: (52) 8248-0101; email: fzavala@ccr.dsi.uanl.mx

2 Department of Agronomy. University of Nebraska–Lincoln. Lincoln, NE. USA. Fax: (402) 472-3066; email: jeastin@unlnotes.unl.edu

3 Facultad de Ciencias Biologicas. UANL. Monterrey, N.L. Mexico. Fax: (52) 8352-1142; email: hgamez@ccr.dsi.uanl.mx

Keywords: respiration, adaptation, sorghum

Two sorghum genotypes (Pioneer 8244® and H2183210351) with different degree of adaptation to tropical areas were planted in march in Marin, N.L. Mexico (tropical environment). The objective was to relate dark respiration with the lack of genetic adaptation. The dark respiration measurements were made 85 days after planting, during the grain fill period. The plants in one meter long of the central row were used for the measurements. A plexiglass chamber of 1 m long, 0.8 m width and 1 m tall was seated on top of a wood base to cover all plants in the meter of the row. The wood base was to avoid CO₂ influx from the soil. The CO₂ measurements inside the camber were made with an infrared gas analyzer LI-COR 6200 (LICOR, CO. USA) at 9:00 hr PM and 05:00 hr AM. The temperature difference between 9:00 PM and 5 AM was 2.3° C. Pioneer was more sensitive in respiration to temperature changes than H2183210351 (3.8 µMolCO₂Kg⁻¹s⁻¹ and 0.7 µMolCO₂Kg⁻¹s⁻¹, respectively). The respiration rate in Pioneer 8244 was 200 % higher than H2183210351 (13.7 µMolCO₂Kg⁻¹s⁻¹ and 4.15 µMolCO₂Kg⁻¹s⁻¹, respectively). The differences in respiration rates are probably due to the lack of adaptation of Pioneer 8244, that it is recommended for temperate environments, compared with the hybrid H2183210351, which parents are more tropical adapted.